

☆ Indexed Journal
☆ Refereed Journal
www.entomoljournal.com

Online-ISSN: 2320-7078
Print-ISSN: 2349-6800
Peer Reviewed Journal

Journal of Entomology and Zoology Studies

Volume 4

Issue 4

Jul-Aug

2016



New Delhi,
India



JOURNAL OF ENTOMOLOGY AND ZOOLOGY STUDIES

(ISSN Print: 2320-7078)

(ISSN Online: 2349-6800)

Volume- 4

Issue- 4


Jul-Aug 2016

Journal of Entomology and Zoology Studies

169, C-11, Sector-3, Rohini, Delhi-110085

Toll Free No.: 18001234070

Mob.: +91-9711224068



THOMSON REUTERS



Year: 2016, Volume: 4, Issue: 4
E-ISSN: 2320-7078
P-ISSN: 2349-6800

Online Available at www.entomoljournal.com



Journal of Entomology and Zoology Studies

Index for 2016 (Vol-4, Issue-4)

Part I

1. **Beekeeping status in Kangra district of Himachal Pradesh**
Rajesh Kumar and Nisha Kundal
Page: 620-622
2. **The Species Composition of Scales and of Armored Scales (Hemiptera, Coccoidea) in Azerbaijan**
Mustafayeva Gulzar Aligeydar
Page: 623-625
3. **Update of distribution records of *Phaonia* Robineau-Desvoidy (Diptera: Muscidae) from Bulgaria with the description of a new species**
Eberhard Zielke
Page: 626-632
4. **Effects of silver nanoparticles (AgNPs) and polymers on stored pests for improving the industry of packaging foodstuffs**
Somaye Allahvaysi
Page: 633-640
5. **Present status of socio-economic conditions of the fishing community of the Meghna River adjacent to Narsingdi District, Bangladesh**
Md. Simul Bhuyan, Aysha Akhtar, KM Saim and Md. Shafiqul Islam
Page: 641-646
6. **First report of four Curculionidae (Coleoptera) from center and northeast of Iran**
Mehdi Zare Khormizi, Minoo Heidari Latibari, Fatemeh Khandehroo, Gholamhossein Moravvej and Hussein Sadeghi Namaghi
Page: 647-649
7. **Influence of food plants on different developmental stages of *Aiolopus thalassinus thalassinus* with special reference to its mating strategies**
Fakhra Soomro, Riffat Sultana and Muhammad Saeed Wagan
Page: 650-653
8. **A New record of Genus *Macrobilharzia* Travassos, 1922 (Trematoda: Schistosomatoidea) in *Phalacrocorax niger* from Sindh, Pakistan**
Muhammad Moosa Abro, Ali Murtaza Dharejo, Muhammad Munif Khan and Nadir Ali Birmani
Page: 654-656
9. **Cotton Mealybug Management: A Review**
Nadia Noreen, Mubashar Hussain, Samman Fatima and Mobein Ghazanfar
Page: 657-663



E-ISSN: 2320-7078
P-ISSN: 2349-6800
JEZS 2016; 4(4): 623-625
© 2016 JEZS
Received: 21-05-2016
Accepted: 22-06-2016

Mustafayeva Gulzar Aligeydar
PhD, Institute of Zoology,
Azerbaijan National Academy of
Sciences, Baku c, Azerbaijan

The Species Composition of Scales and of Armored Scales (Hemiptera, Coccoidea) in Azerbaijan

Mustafayeva Gulzar Aligeydar

Abstract

The article presents faunistical research data on both scales and armored scales that are wide spread in Azerbaijan. Among the whole number of those identified, 52 species of Coccoidea including 11 species of scales (Coccoidea, Diaspididae) and 8 species of armored scales (Coccoidea, Lecaniidae) were found for the first time in Azerbaijan's fauna.

Keywords: pest, scales, armored scales, fauna, species composition, new species

Introduction

The representatives of the order Hemiptera (Homoptera) have a special place among the pests of agricultural crops and of decorative plantings. Both scales and armored scales (Coccoidea), which are members of that order, make big damages to agricultural crops and to decorative plants. They suck out juices from the plant there by causing desiccation, atrophy and abscission of leaves, dry down of branches, distortion of leaves, fruits and offshoots, and reduction of the annual amount of growth in plants.

They seriously aggravate condition of agricultural crops, as well as greatly reduce both quality and quantity of the harvest sometimes even leading to its complete loss. The decorative and forest-park culture affected by the mentioned invaders lose their beauty, and when a mass infestation comes, they dry out completely. The mentioned pests reproduce with significant speed causing great damage to plants and leading to their death on occasions. Many of those species are polyphagous with high fertility rate and with ecological plasticity, which allow them to spread out widely and to cause much harm to the plants.

There were very little studies done on the fauna of scales and armored scales of Azerbaijan. The original information came from the research of E.S. Arutyunova and V. Kh. Rusanova^[1].^[7] A. Imamkuliyeve reported some data on scales and armored scales affecting fruit trees and subtropical cultures in the Lankaran area of Azerbaijan^[5, 6]. R.A. Beybutov presented information about parasites of brown soft scale wide spread in Azerbaijan^[2, 3]. L.M. Rzaeva mentions in her monograph some species of scales and armored scales common in Azerbaijan^[18, 19]. G.A. Mustafayeva in her publications writes about scales and armored scales as hosts of some species of Aphelinidae^[7-10]. Same author reports about certain biological peculiarities of both Japanese wax scale and globose scale, as well as about scales of decorative crops^[11-13]. The works of G. A. Mustafayeva, V. P. Kamarli, G. M. Aslanova, B.A. Ahmedov, D.V. Mustafayev, N.A. Hasanov, and S.B. Ahmedov specify data related to fauna of armored scales and scales of Guba-Khachmaz area in Azerbaijan^[14, 15]. G. A. Mustafayeva, I. Karaca, G.J. Statas, M.S. Ozgekce and P.J. Skouras discuss the fauna of scales harming park and decorative cultures in Azerbaijan, Turkey and Greece^[16].

Material and methods

The entomological material has been collected from natural and cultural biocoenoses using common methodology (Borchsenius, 1950; Shapiro, Tryapitsyn, Shepetilnikova, 1982)^[4, 20]. The pests have been collected during individual and complex faunistical expeditions and research trips. The preparation and development of entomological material has been done in the Institute of Zoology of Azerbaijan National Academy of Sciences, namely in the Laboratory of Introduction of Useful Insects and Principles of Biological Control. Both scales

Correspondence

Mustafayeva Gulzar Aligeydar
PhD, Institute of Zoology,
Azerbaijan National Academy of
Sciences, Baku c, Azerbaijan

and armored scales have been stored with excised parts of the plants. In order to determine the plant species along with the pest, there were prepared herbaria from different types of plants, after which the specific Latin name of each plant was identified.

Results and Discussion

As a result of the number of years of faunistic research in Azerbaijan, there have been detected 33 species of scales belonging to 21 families [7-9]. That number has included 11 species found in Azerbaijani fauna for the first time:

Lepidosaphes granati (Koroneos), *Aulacaspis rosae* (Bouche), *Pseudaulacaspis pentagona* (Targioni-Tozzetti), *Carulaspis visci* (Schränk.), *Carulaspis minima* (Targioni-Tozzetti), *Adiscodiaspis tamaricicola* Malenotti, *Diaspidiotus caucasicus* (Borchsenius), *Tecaspis prunorum* (Borchsenius), *Tecaspis asiatica* Balachowsky, *Aonidia lauri* Bouche., *Lopholeucaspis yaponica* Cock [10, 11].

Both systematic list and species composition of scale insects is given in Table 1 below. The new species of scales for Azerbaijan are marked with an asterisk (*).

Table 1: Generic and species composition of scales (Hemiptera, Diaspididae) in Azerbaijan

Generes of scales	Species of scales
1. <i>Parlatoria</i> Targioni – Tozzetti, 1868	<i>Parlatoria oleae</i> (Colvée, 1880) <i>Parlatoria ziziphi</i> (Lucas, 1853)
2. <i>Leucaspis</i> Targioni – Tozzetti, 1868	<i>Leucaspis pusilla</i> Low., 1883
3. <i>Lepidosaphes</i> Shimer L., 1868	<i>Lepidosaphes ulmi</i> (Linnaeus, 1758) <i>Lepidosaphes gloveri</i> (Packard, 1869) <i>Lepidosaphes ficus</i> Sign., 1870 <i>Lepidosaphes granati</i> (Koroneos, 1934)*
4. <i>Cornuaspis</i> Mac Gillivray, 1921.	<i>Cornuaspis beckeri</i> (Newman, 1869)
5. <i>Unaspis</i> Mac Gillivray, 1921.	<i>Unaspis evonymi</i> (Comstock, 1881)
6. <i>Aulacaspis</i> Cockerell, 1893.	<i>Aulacaspis rosae</i> (Bouche, 1833) *
7. <i>Pseudaulacaspis</i> Mac Gillivray, 1921.	<i>Pseudaulacaspis pentagona</i> (Targioni-Tozzetti, 1885) *
8. <i>Carulaspis</i> Mac Gillivray, 1921.	<i>Carulaspis visci</i> (Schränk., 1781) * <i>Carulaspis minima</i> (Targioni-Tozzetti, 1868) *
9. <i>Adiscodiaspis</i> Marchal, 1909.	<i>Adiscodiaspis tamaricicola</i> Malenotti, 1916*
10. <i>Chrysomphalus</i> Ashmead, 1880.	<i>Chrysomphalus dictyospermi</i> (Morgan, 1889)
11. <i>Kuwanaspis</i> Mac Gillivray, 1921.	<i>Kuwanaspis howardi</i> (Cooley, 1898)
12. <i>Aonidiella</i> Berlese et Leonardi, 1895	<i>Aonidiella citrine</i> (Coquillett, 1891)
13. <i>Aspidiotus</i> Bouche, 1833.	<i>Aspidiotus nerii</i> Bouche, 1937
14. <i>Temnaspis</i> Signoret, 1869	<i>Temnaspis destructor</i> (Signoret, 1869)
15. <i>Epidiaspis</i> Cockerell, 1899	<i>Epidiaspis leperii</i> Signoret, 1869
16. <i>Tecaspis</i> Hall., 1929.	<i>Tecaspis prunorum</i> (Borchsenius, 1939) * <i>Tecaspis asiatica</i> Balachowsky, 1954*
17. <i>Diaspidiotus</i> Berlet Leon, 1895.	<i>Diaspidiotus slavonicus</i> (Qrenn., 1934) <i>Diaspidiotus pyri</i> (Lichtenstein, 1881) <i>Diaspidiotus gigas</i> (Theim and Gerneck., 1934) <i>Diaspidiotus ostreaformis</i> (Curtis, 1843) <i>Diaspidiotus caucasicus</i> (Borchsenius, 1935)* <i>Diaspidiotus perniciosus</i> (Comstock, 1881) <i>Diaspidiotus prunorum</i> (Laing., 1931)
18. <i>Salicicola</i> Lindinger, 1905.	<i>Salicicola kermanensis</i> (Lindinger, 1905.)
19. <i>Diaspis</i> Costa, 1828.	<i>Diaspis echinocacti</i> (Bouche, 1933)
20. <i>Lopholeucaspis</i> Cockerell, 1897	<i>Lopholeucaspis yaponica</i> Balach., 1953*
21. <i>Aonidia</i> Targioni-Tozzetti, 1868.	<i>Aonidia lauri</i> (Bouche, 1833)*

Based on the number of years of research of Azerbaijani fauna, there have been also detected 19 species of armored scales belonging to nine families [7, 8]. Of that number, eight species have been discovered in Azerbaijan's fauna for the first time: *Eulecanium rufulum* Ckll., *Eulecanium rugulosum* (Arc.), *Pulvinaria aurantii*, Ckll., *Pulvinaria artemisiae* Sign.,

Pulvinaria pistaciae Bod., *Saissetia oleae* (Bern.), *Ceroplastes sinensis* Guer., *Physokermes piceae* (Sch.). Those species are marked with two asterisks (**).

Both systematic list and species composition of scale insects is given in Table 2 below.

Table 2: Generic and species composition of armored scales (Hemiptera, Lecanidae) in Azerbaijan

Generes of armored scales	Species of armored scales
1. <i>Coccus</i> L., 1758.	<i>Coccus hesperidum</i> L. 1758 <i>Coccus pseudomagnoliarum</i> (Kuw., 1914)
2. <i>Eulecanium</i> Ckll., 1896	<i>Eulecanium bituberculatum</i> Targ., 1869 <i>Eulecanium rufulum</i> Ckll., 1903** <i>Eulecanium rugulosum</i> (Arc, 1937)**
3. <i>Parthenolecanium</i> Sulc., 1908	<i>Parthenolecanium corni</i> (Bouche, 1844) <i>Parthenolecanium persicae</i> (Fabricius, 1776)
4. <i>Sphaerolecanium</i> Sulc., 1908	<i>Sphaerolecanium prunastri</i> Fonscolombe, 1873
5. <i>Rhodococcus</i> Borchs., 1953.	<i>Rhodococcus turanicus</i> (Arch., 1937)
6. <i>Pulvinaria</i> Targioni Tozzetti, 1869	<i>Pulvinaria betulae</i> (L. 1758) <i>Pulvinaria aurantii</i> Ckll., 1896 **

	<i>Pulvinaria floccifera</i> (Westw., 1870) <i>Pulvinaria artemisiae</i> Sign, 1873 ** <i>Pulvinaria pistaciae</i> Bod, 1926**
7. <i>Saissetia</i> Depl, 1865.	<i>Saissetia oleae</i> (Bern, 1782)** <i>Saissetia hemispaerica</i> (Targ., 1867)
8. <i>Ceroplastes</i> Gray, 1830.	<i>Ceroplastes yaponicus</i> Green, 1921. <i>Ceroplastes sinensis</i> Guer., 1900.**
9. <i>Physokermes</i> Targ., 1869.	<i>Physokermes piceae</i> (Sch., 1801)**

Conclusions

1. As a result of the unique and complex study of Hemiptera (Coccoidea) in Azerbaijan, there were detected 52 species of both scales and armored scales. 33 species among them were found to be scales (Coccoidea, Diaspididae) which belong to 21 genus. The remaining 19 species were found to be armored scales (Coccoidea, Lecaniidae) which belong to nine genera.

2. Among all of the 33 identified species of scale insects (Hemiptera, Diaspididae), 11 species were found in Azerbaijan's fauna for the first time. The genus *Diaspidiotus* is represented by seven species, and the genus *Lepidosaphes* is represented by four species. The genera *Parlatoria*, *Carulaspis* and *Tecaspis* are represented by two species in each of them respectively. Each of the 16 genera has only one species. Majority of species discovered were polyphagous, eight species were monophagous, and six species were oligophagous.

3. Among all of the 19 identified species of armored scales (Hemiptera, Lecaniidae) eight species were found in Azerbaijan's fauna for the first time. The genus *Pulvinaria* is represented by five species, the genus *Eulecanium* is represented by three species. The genera *Coccus*, *Parthenolecanium*, *Saissetia* and *Ceroplastes* are represented by two species in each of them respectively. Each of the genera *Rhodococcus* and *Physokermes* has only one species in them. Only *Pulvinaria pistaciae*, *Pulvinaria artemisiae* and *Physokermes piceae* species were monophagous while the remaining species were either oligophagous or polyphagous.

References

- Arutyunova ES. The overview of fauna of coccids of Azerbaijan. Baku, Azerbaijan, 1938, 24-54.
- Beybutov RA. The integrated method of control of brown soft scales. Reports Azerbaijan SSR, 1956; 12(9):661-664.
- Beybutov RA. The biological method of pest control on perennial crops in Azerbaijan conditions, Collection Az. SSR, dedicated to XXI Congress of CPSU, Baku, Azerbaijan, 1960, 349-353.
- Borchsenius NS. The practical manual of coccids (Coccoidea) of cultivated plants and forest species of USSR. Moscow--Leningrad, USSR, 1950, 27-75.
- Imamkuliyevev AG. The coccids (Homoptera, Coccoidea) harmful for fruit and subtropical cultures in Lankaran area of Azerbaijan. Bulletin of Azerb. AS, 1966; 4:45-51.
- Imamkuliyevev AG. The coccids and the scale insects (Homoptera, Coccidae) harmful for fruit and subtropical cultures in Lankaran area of Azerbaijan. Abstract of Cand.Sc. (Biol.) dissertation. Baku, Azerbaijan, 1966, 1-20.
- Mustafayeva GA. The aphelinids (Hymenoptera, Aphelinidae) of the Eastern Azerbaijan (fauna, ecology and economical importance). Abstract of Cand. Sc. (Biol.) dissertation. 1990, 1-20.
- Mustafayeva GA. The aphelinids (Hymenoptera, Aphelinidae) as parasites of coccids, aphids and whiteflies in Eastern Azerbaijan. Bulletin of Azerb. AS. 2004, 1-2:91-101.
- Mustafayeva GA. Some species of scales (Hemiptera, Diaspididae) harmful to decorative cultures of Absheron peninsula and their entomophagous. Proceedings of International Scientific Conference on "The Introduction and Protection of Plants in Botanical Gardens and Arboretums. 2009, 108-113.
- Mustafayeva GA. The aphelinids (Hymenoptera, Aphelinidae) as parasites of sucking pests of Azerbaijan. Proceedings of VI International Scientific Conference on Industrial Botany. Donetsk, Ukraine, 2010, 318-320.
- Mustafayeva GA. The fauna of scales (Hemiptera, Diaspididae) of Azerbaijan; their occurrence and harmfulness. Proceedings of Azerbaijan Society of Zoologists. Baku, Azerbaijan, 2013, 65-77.
- Mustafayeva GA. Japanese wax scale (*Ceroplastes yaponicus* Green.) and globose scale (*Sphaerolecanium prunastri* Fonsc.) with their the entomophagous in North-Eastern Azerbaijan. The journal "Current issues in humanities and natural sciences. Moscow, Russia, 2015; 5:31-36.
- Mustafayeva GA. On three species of scales (Hemiptera, Diaspididae) damaging the ornamental plants in Azerbaijan and their entomophagous. Proceedings of the 7th International Conference on European Science and Technology. April 23 - 24, Munich, Germany, 2014, 99-03.
- Mustafayeva GA, Ahmedov BA, Kamarli VP, Aslanova GM, Mustafayev DV et al. The scales (Hemiptera: Diaspididae) of Guba-Khachmaz area of Azerbaijan; their distribution and trophic relationship. The International Conference on European Science and Technology, 24-25-2014, Munich, Germany, 2014, 69-73.
- Mustafayeva GA, Kamarli VP, Aslanova GM. The armored scales (Homoptera, Coccinea, Lecaniidae) of Guba-Khachmaz area of Azerbaijan, their distribution and trophic communications. Austrian Journal of Technical and Natural Sciences Vienna Austria. 2014, 11-12:10-13.
- Mustafayeva GA, Karaca I, Stathas GJ, Ozgokce MS, Skouras PJ. On some scales (Hemiptera, Diaspididae) infesting ornamental and forest park plants in Azerbaijan, Greece and Turkey. III International Scientific Conference The Introduction and Protection of Plants. 2012, 172.
- Rusanova VN. Materials to study Coccidae (Homoptera) in Azerbaijan. Baku, Azerbaijan, 1941, 4-20.
- Rzayeva LM. The chalcids (Hymenoptera, Chalcidoidea) of Eastern Transcaucasia and their economical value. Abstract of PhD dissertation. Baku, Azerbaijan, 1987, 1-45.
- Rzayeva LM. The chalcids (Hymenoptera, Chalcidoidea) of Eastern Transcaucasia and their economic value. Baku, Azerbaijan, 2002, 1-354.
- Tryapitsyn VA, Shapiro VA, Shepetilnikova VA. Parasites and predators of pests of agricultural crops. Leningrad, USSR: Kolos, 1982, 256.



E-ISSN: 2320-7078
P-ISSN: 2349-6800



Journal of Entomology and Zoology Studies

Publication Certificate

This certificate confirms that "Mustafayeva Gulzar Aligeydar", has published manuscript titled "The Species Composition of Scales and of Armored Scales (Hemiptera, Coccoidea) in Azerbaijan".

Details of Published Article as follow

Volume	:	4
Issue	:	4
Year	:	2016
Page Number	:	623-625
Article Regd. No	:	Ento-4-4-173

Certificate No.: 4-4-173
Date: 29-07-2016

Yours Sincerely,

Akhil Gupta

Akhil Gupta
Publisher
Journal of Entomology and Zoology Studies
Website: www.entomoljournal.com

